

# Approximating Maximum Edge 2-Coloring in Simple Graphs

Zhi-Zhong Chen\*

Sayuri Konno<sup>†</sup>

Yuki Matsushita<sup>‡</sup>

## Abstract

We present a polynomial-time approximation algorithm for legally coloring as many edges of a given simple graph as possible using two colors. It achieves an approximation ratio of roughly 0.842 and runs in  $O(n^3m)$  time, where  $n$  (respectively,  $m$ ) is the number of vertices (respectively, edges) in the input graph. The previously best ratio achieved by a polynomial-time approximation algorithm was  $\frac{5}{6} \approx 0.833$ .

**Keywords:** Approximation algorithms, graph algorithms, edge coloring, NP-hardness.

## 1 Introduction

Given a graph  $G$  and a natural number  $t$ , the *maximum edge  $t$ -coloring problem* (called MAX EDGE  $t$ -COLORING for short) is to find a maximum-sized set  $F$  of edges in  $G$  such that  $F$  can be partitioned into at most  $t$  matchings of  $G$ . Motivated by call admittance issues in satellite based telecommunication networks, Feige et al. [3] introduced the problem and proved its APX-hardness. They also observed that MAX EDGE  $t$ -COLORING is a special case of the well-known maximum coverage problem (see [6]). Since the maximum coverage problem can be approximated by a greedy algorithm within a ratio of  $1 - (1 - \frac{1}{t})^t$  [6], so can MAX EDGE  $t$ -COLORING. In particular, the greedy algorithm achieves an approximation ratio of  $\frac{3}{4}$  for MAX EDGE 2-COLORING, which is the special case of MAX EDGE  $t$ -COLORING where the input number  $t$  is fixed to 2. For this special case, Feige et al. [3] has improved the trivial ratio  $\frac{3}{4} = 0.75$  to  $\frac{10}{13} \approx 0.769$  by an LP approach.

The APX-hardness proof for MAX EDGE  $t$ -COLORING given by Feige et al. [3] indeed shows that the problem remains APX-hard even if we restrict the input graph to a simple graph and fix the input integer  $t$  to 2. We call this restriction (special case) of the problem MAX SIMPLE EDGE 2-COLORING. Feige et al. [3] also pointed out that for MAX SIMPLE EDGE 2-COLORING, an approximation ratio of  $\frac{4}{5}$  can be achieved by the following *simple algorithm*: Given a simple graph  $G$ , first compute a maximum-sized subgraph  $H$  of  $G$  such that the degree of each vertex in  $H$  is at most 2 and there is no 3-cycle in  $H$ , and then remove one *arbitrary* edge from each odd cycle

---

\*Corresponding author. Supported in part by the Grant-in-Aid for Scientific Research of the Ministry of Education, Science, Sports and Culture of Japan, under Grant No. 20500021. Department of Mathematical Sciences, Tokyo Denki University, Hatoyama, Saitama 350-0394, Japan. Email: zzchen@mail.dendai.ac.jp

<sup>†</sup>Department of Mathematical Sciences, Tokyo Denki University, Hatoyama, Saitama 350-0394, Japan.

<sup>‡</sup>Department of Mathematical Sciences, Tokyo Denki University, Hatoyama, Saitama 350-0394, Japan.

of  $H$ . This simple algorithm has been improved in [1, 2, 9]. The previously best ratio (namely,  $\frac{5}{6}$ ) was given in [9]. In this paper, we improve on both the algorithm in [1] and the algorithm in [9] to obtain a new approximation algorithm that achieves a ratio of roughly 0.842. Roughly speaking, our algorithm is based on local improvement, dynamic programming, and recursion. Its analysis is based on an intriguing charging scheme and certain structural properties of *train* graphs and *starlike* graphs (see Section 3 for definitions).

Kosowski et al. [10] also considered MAX SIMPLE EDGE 2-COLORING. They presented an approximation algorithm that achieves a ratio of  $\frac{28\Delta-12}{35\Delta-21}$ , where  $\Delta$  is the maximum degree of a vertex in the input simple graph. This ratio can be arbitrarily close to the trivial ratio  $\frac{4}{5}$  because  $\Delta$  can be very large. In particular, this ratio is worse than our new ratio 0.842 when  $\Delta \geq 4$ . Moreover, when  $\Delta = 3$ , our algorithm indeed achieves a ratio of  $\frac{6}{7}$ , which is equal to the ratio  $\frac{28\Delta-12}{35\Delta-21}$  achieved by Kosowski et al.'s algorithm [10]. Note that MAX SIMPLE EDGE 2-COLORING becomes trivial when  $\Delta \leq 2$ . Therefore, no matter what  $\Delta$  is, our algorithm is better than or as good as all known approximation algorithms for MAX SIMPLE EDGE 2-COLORING.

Kosowski et al. [10] showed that approximation algorithms for MAX SIMPLE EDGE 2-COLORING can be used to obtain approximation algorithms for certain packing problems and fault-tolerant guarding problems. Combining their reductions and our improved approximation algorithm for MAX SIMPLE EDGE 2-COLORING, we can obtain improved approximation algorithms for their packing problems and fault-tolerant guarding problems immediately.

## 2 Basic Definitions

Throughout the remainder of this paper, a graph means a simple undirected graph (i.e., it has neither parallel edges nor self-loops).

Let  $G$  be a graph. We denote the vertex set of  $G$  by  $V(G)$ , and denote the edge set of  $G$  by  $E(G)$ . The *degree* of a vertex  $v$  in  $G$ , denoted by  $d_G(v)$ , is the number of vertices adjacent to  $v$  in  $G$ . A vertex  $v$  of  $G$  with  $d_G(v) = 0$  is called an *isolated vertex*. For a subset  $U$  of  $V(G)$ , let  $G[U]$  denote the graph  $(U, E_U)$  where  $E_U$  consists of all edges  $\{u, v\}$  of  $G$  with  $u \in U$  and  $v \in U$ . We call  $G[U]$  the *subgraph of  $G$  induced by  $U$* . For a subset  $U$  of  $V(G)$ , we use  $G - U$  to denote  $G[V(G) - U]$ .  $G$  is a *star* if  $G$  is connected,  $G$  has at least three vertices, and there is a vertex  $u$  (called the *center* of  $G$ ) such that every edge of  $G$  is incident to  $u$ . Each vertex of a star other than the center is called a *satellite* of the star.

A *cycle* in  $G$  is a connected subgraph of  $G$  in which each vertex is of degree 2. A *path* in  $G$  is a connected subgraph of  $G$  in which exactly two vertices are of degree 1 and the others are of degree 2. Each vertex of degree 1 in a path  $P$  is called an *endpoint* of  $P$ , while each vertex of degree 2 in  $P$  is called an *inner vertex* of  $P$ . An edge  $\{u, v\}$  of a path  $P$  is called an *inner edge* of  $P$  if both  $u$  and  $v$  are inner vertices of  $P$ . The *length* of a cycle or path  $C$  is the number of edges in  $C$ . A cycle of odd (respectively, even) length is called an *odd* (respectively, *even*) cycle.

A *path-cycle cover* of  $G$  is a subgraph  $H$  of  $G$  such that  $V(H) = V(G)$  and  $d_H(v) \leq 2$  for every  $v \in V(H)$ . Note that each connected component of a path-cycle cover of  $G$  is an isolated vertex,

path, or cycle. A path-cycle cover  $\mathcal{C}$  of  $G$  is *triangle-free* if  $\mathcal{C}$  does not contain a cycle of length 3. A path-cycle cover  $C$  of  $G$  is *maximum-sized* if the number of edges in  $C$  is maximized over all path-cycle covers of  $G$ .

$G$  is *edge-2-colorable* if each connected component of  $G$  is an isolated vertex, a path, or an even cycle. Note that MAX SIMPLE EDGE 2-COLORING is the problem of finding a maximum-sized edge-2-colorable subgraph in a given graph.

### 3 Two Crucial Lemmas and the Outline of Our Algorithm

We say that a graph  $K = (V_K, E_K \cup F_K)$  is a *train graph* if it satisfies the following conditions:

- The graph  $(V_K, E_K)$  has  $h + 1$  connected components  $C_0, \dots, C_h$  with  $h \geq 0$ .
- $C_0$  is a path while  $C_1$  through  $C_h$  are odd cycles of length at least 5.
- $F_K$  is a matching consisting of  $h$  edges  $\{u_1, v_1\}, \dots, \{u_h, v_h\}$ .
- For each  $i \in \{1, \dots, h\}$ ,  $u_i$  is an inner vertex of path  $C_0$  while  $v_i$  is a vertex of  $C_i$ .

We call the edges of  $F_K$  the *column edges* of  $K$ , call path  $C_0$  the *beam path* of  $K$ , and call cycles  $C_1$  through  $C_h$  the *wheels* of  $K$ .

We say that a graph  $K = (V_K, E_K \cup F_K)$  is a *starlike graph* if it satisfies the following conditions:

- The graph  $(V_K, E_K)$  has  $h + 1$  connected components  $C_0, \dots, C_h$  with  $h \geq 2$ .
- $C_0$  is a cycle of length at least 4 while  $C_1$  through  $C_h$  are odd cycles of length at least 5.
- $F_K$  is a matching consisting of  $h$  edges  $\{u_1, v_1\}, \dots, \{u_h, v_h\}$ .
- For each  $i \in \{1, \dots, h\}$ ,  $u_i$  is a vertex of  $C_0$  while  $v_i$  is a vertex of  $C_i$ .

We call the edges of  $F_K$  the *bridge edges* of  $K$ , call  $C_0$  the *central cycle* of  $K$ , and call  $C_1$  through  $C_h$  the *satellite cycles* of  $K$ .

Let  $r$  be the root of the quadratic equation  $23r^2 - 55r + 30 = 0$  that is smaller than 1. Note that  $r = 0.84176\dots \approx 0.842$ . The reason why we choose  $r$  in this way will become clear later in the proof of Lemma 4.8.

**Lemma 3.1** *Suppose that  $K$  is a train graph such that each wheel of  $K$  is charged a penalty of  $6 - 7r$ . Let  $p(K)$  be the total penalties charged to the wheels of  $K$ . Then,  $K$  has an edge-2-colorable subgraph  $K'$  such that  $|E(K')| - p(K) \geq r|E_K|$ , where  $E_K$  is the set of edges on the beam path or the wheels of  $K$ .*

PROOF. Note that the degree of each vertex in  $K$  is at most 3. We prove the lemma by induction on  $\kappa$  which is the number of edges  $e$  on the beam path of  $K$  such that both endpoints of  $e$  are of degree 3 in  $K$ .

(Basis) In the base case,  $\kappa = 0$ . We obtain  $K'$  from  $K$  by deleting the column edges of  $K$  and removing one edge from each wheel of  $K$ . Let  $\tau$  be the number of wheels of  $K$ . Then,  $|E(K')| = |E_K| - \tau$ . Moreover,  $|E_K| \geq 5\tau + 2\tau = 7\tau$  because the wheels of  $K$  contain at least  $5\tau$  edges while the beam path of  $K$  contains at least  $2\tau$  edges for  $\kappa = 0$ . So,  $|E(K')| - p(K) \geq r|E_K|$  because  $p(K) = (6 - 7r)\tau$ .

(Induction step) Suppose that  $\kappa \geq 1$ . Let  $\{u_1, u_2\}$  be an arbitrary edge on the beam path of  $K$  such that both  $u_1$  and  $u_2$  are of degree 3 in  $K$ . For each  $i \in \{1, 2\}$ , let  $\{u_i, v_i\}$  be the column edge of  $K$  incident to  $u_i$ , and let  $C_i$  be the wheel of  $K$  with  $v_i \in V(C_i)$ . We cut  $K$  into two train graphs  $K_1$  and  $K_2$  by deleting edge  $\{u_1, u_2\}$ , deleting column edges  $\{u_1, v_1\}$  and  $\{u_2, v_2\}$ , and deleting wheels  $C_1$  and  $C_2$ . By the inductive hypothesis,  $K_i$  has an edge-2-colorable subgraph  $K'_i$  with  $|E(K'_i)| - p(K_i) \geq r|E(K_i) \cap E_K|$  for each  $i \in \{1, 2\}$ , where  $p(K_i)$  is the total penalties charged to the wheels of  $K_i$ . We obtain  $K'$  from  $K'_1$  and  $K'_2$  by adding column edges  $\{u_1, v_1\}$  and  $\{u_2, v_2\}$ , the path obtained from  $C_1$  by removing one edge incident to  $v_1$ , and the path obtained from  $C_2$  by removing one edge incident to  $v_2$ . Clearly,  $|E(K')| = \sum_{i=1}^2 (|E(K'_i)| + |E(C_i)|)$ . So,  $|E(K')| - \sum_{i=1}^2 p(K_i) \geq \sum_{i=1}^2 (r|E(K_i) \cap E_K| + |E(C_i)|)$ . Note that  $p(K) = \sum_{i=1}^2 p(K_i) + 2(6 - 7r)$  and  $|E_K| = \sum_{i=1}^2 (|E(K_i) \cap E_K| + |E(C_i)|) + 1$ . Now, since  $|E(C_1)| + |E(C_2)| \geq 10$  and  $r \geq \frac{3}{4}$ , we have  $|E(K')| - p(K) \geq r|E_K|$ .  $\square$

**Lemma 3.2** *Suppose that  $K$  is a starlike graph such that each satellite cycle of  $K$  is charged a penalty of  $6 - 7r$ . Let  $p(K)$  be the total penalties charged to the satellite cycles of  $K$ . Then,  $K$  has an edge-2-colorable subgraph  $K'$  such that  $|E(K')| - p(K) \geq r|E_K|$ , where  $E_K$  is the set of edges on the central or satellite cycles of  $K$ .*

PROOF. Let  $C_0$  be the central cycle of  $K$ . Let  $C_1, \dots, C_h$  be the satellite cycles of  $K$ . We distinguish two cases as follows.

*Case 1:* No two degree-3 vertices are adjacent in  $K$ . In this case, we obtain  $K'$  from  $K$  as follows:

- For every  $i \in \{2, \dots, h\}$ , remove one edge of  $C_i$  incident to the bridge edge between  $C_0$  and  $C_i$ , and further remove the bridge edge.
- For the bridge edge  $\{u_0, u_1\}$  between  $C_0$  and  $C_1$  with  $u_0 \in V(C_0)$  and  $u_1 \in V(C_1)$ , remove one edge of  $C_0$  incident to  $u_0$  and remove one edge of  $C_1$  incident to  $u_1$ .

Clearly,  $|E(K')| = \sum_{i=0}^h |E(C_i)| - h$  and  $p(K) = (6 - 7r)h$ . Moreover, since  $|E(C_0)| \geq 2h$  and  $|E(C_i)| \geq 5$  for each  $i \in \{1, \dots, h\}$ , we have  $|E_K| = \sum_{i=0}^h |E(C_i)| \geq 7h$ . Thus,  $|E(K')| - p(K) \geq r|E_K|$ .

*Case 2:* There are two degree-3 vertices adjacent in  $K$ . In this case, there is an edge  $\{u_1, u_2\} \in E(C_0)$  such that both  $u_1$  and  $u_2$  are of degree 3 in  $K$ . Without loss of generality, we may assume that for each  $i \in \{1, 2\}$ ,  $C_i$  contains the vertex  $v_i$  such that  $\{u_i, v_i\}$  is the bridge edge of  $K$  between  $C_0$  and  $C_i$ . Consider the train  $K_1$  obtained from  $K$  by deleting edge  $\{u_1, u_2\}$ , deleting bridge edges  $\{u_1, v_1\}$  and  $\{u_2, v_2\}$ , and deleting satellite cycles  $C_1$  and  $C_2$ . By Lemma 3.1, we can obtain

an edge-2-colorable subgraph  $K'_1$  of  $K_1$  with  $|E(K'_1)| - p(K_1) \geq r|E(K_1) \cap E_K|$ . We obtain  $K'$  from  $K'_1$  by adding bridge edges  $\{u_1, v_1\}$  and  $\{u_2, v_2\}$ , the path obtained from  $C_1$  by removing one edge incident to  $v_1$ , and the path obtained from  $C_2$  by removing one edge incident to  $v_2$ . Clearly,  $|E(K')| = |E(K'_1)| + |E(C_1)| + |E(C_2)|$ . So,  $|E(K')| - p(K_1) \geq r|E(K_1) \cap E_K| + |E(C_1)| + |E(C_2)|$ . Note that  $|E_K| = |E(K_1) \cap E_K| + |E(C_1)| + |E(C_2)| + 1$  and  $p(K) = p(K_1) + 2(6 - 7r)$ . Now, since  $\sum_{i=1}^2 |E(C_i)| \geq 10$  and  $r \geq \frac{2}{3}$ , we have  $|E(K')| - p(K) \geq r|E_K|$ .  $\square$

Based on Lemmas 3.1 and 3.2, we will design our algorithm roughly as follows: Given an input graph  $G$ , we will first construct a suitable maximum-sized triangle-free path-cycle cover  $\mathcal{C}$  of  $G$  and compute a suitable set  $F$  of edges such that the endpoints of each edge in  $F$  fall into different connected components of  $\mathcal{C}$  and each odd cycle of  $\mathcal{C}$  has at least one vertex that is an endpoint of an edge in  $F$ . Note that  $\mathcal{C}$  has at least as many edges as a maximum-sized edge-2-colorable subgraph of  $G$ . The edges in  $F$  will play the following role: we will break each odd cycle  $C$  in  $\mathcal{C}$  by removing one edge of  $C$  incident to an edge of  $F$  and then this edge of  $F$  can possibly be added to  $\mathcal{C}$  so that  $\mathcal{C}$  becomes an edge-2-colorable subgraph of  $G$ . Unfortunately, not every edge of  $F$  can be added to  $\mathcal{C}$  and we have to discard some edges from  $F$ , leaving some odd cycles of  $\mathcal{C}$   $F$ -free (i.e., having no vertex incident to an edge of  $F$ ). Clearly, breaking an  $F$ -free odd cycle  $C$  of short length (namely, 5) by removing one edge from  $C$  results in a significant loss of edges from  $\mathcal{C}$ . We charge the loss to the non- $F$ -free odd cycles (unevenly) as penalties. Fortunately, adding the edges of  $F$  to  $\mathcal{C}$  will yield a graph whose connected components are train graphs, starlike graphs, or certain other kinds of graphs with good properties. Now, Lemmas 3.1 and 3.2 help us show that our algorithm achieves a ratio of  $r$ .

## 4 The Algorithm

Throughout this section, fix a graph  $G$  and a maximum-sized edge-2-colorable subgraph  $\mathcal{B}$  (for “best”) of  $G$ . Let  $n$  (respectively,  $m$ ) be the number of vertices (respectively, edges) in  $G$ . Our algorithm starts by performing the following four steps:

1. If  $|V(G)| \leq 2$ , then output  $G$  itself and halt.
2. Compute a maximum-sized triangle-free path-cycle cover  $\mathcal{C}$  of  $G$ . (*Comment:* This step can be done in  $O(n^2m)$  time [5].)
3. While there is an edge  $\{u, v\} \in E(G) - E(\mathcal{C})$  such that  $d_{\mathcal{C}}(u) \leq 1$  and  $v$  is a vertex of some cycle  $C$  of  $\mathcal{C}$ , modify  $\mathcal{C}$  by deleting one (arbitrary) edge of  $C$  incident to  $v$  and adding edge  $\{u, v\}$ .
4. Construct a graph  $G_1 = (V(G), E_1)$ , where  $E_1$  is the set of all edges  $\{u, v\} \in E(G) - E(\mathcal{C})$  such that  $u$  and  $v$  appear in different connected components of  $\mathcal{C}$  and at least one of  $u$  and  $v$  appears on an odd cycle of  $\mathcal{C}$ .

Hereafter,  $\mathcal{C}$  always refers to the path-cycle cover obtained after the completion of Step 3. We give several definitions related to the graphs  $G_1$  and  $\mathcal{C}$ . Let  $S$  be a subgraph of  $G_1$ .  $S$  *saturates* an odd cycle  $C$  of  $\mathcal{C}$  if at least one edge of  $S$  is incident to a vertex of  $C$ . The *weight* of  $S$  is the number of odd cycles of  $\mathcal{C}$  saturated by  $S$ . For convenience, we say that two connected components  $C_1$  and  $C_2$  of  $\mathcal{C}$  are *adjacent* in  $G$  if there is an edge  $\{u_1, u_2\} \in E(G)$  such that  $u_1 \in V(C_1)$  and  $u_2 \in V(C_2)$ .

**Lemma 4.1** *We can compute a maximum-weighted path-cycle cover in  $G_1$  in  $O(nm \log n)$  time.*

PROOF. The proof is similar to that of Proposition 2.2 in [9], and is hence by a reduction to the maximum-weight  $[f, g]$ -factor problem. Recall that for two functions  $f$  and  $g$  mapping each vertex  $v$  of a graph  $\mathcal{G}$  to an integer with  $f(v) \leq g(v)$ , an  $[f, g]$ -factor of  $\mathcal{G}$  is a subgraph  $\mathcal{H}$  of  $\mathcal{G}$  such that  $V(\mathcal{H}) = V(\mathcal{G})$  and  $f(v) \leq d_{\mathcal{H}}(v) \leq g(v)$  for every  $v \in V(\mathcal{G})$ . The *weight* of an  $[f, g]$ -factor  $\mathcal{H}$  of  $\mathcal{G}$  is the total weight of edges in  $\mathcal{H}$ . It is known that a maximum-weight  $[f, g]$ -factor of a given edge-weighted graph with  $n'$  vertices and  $m'$  edges can be computed in  $O(n'm' \log n')$  time [4].

Let  $C_1, \dots, C_k$  be the odd cycles of  $\mathcal{C}$ . We construct an auxiliary edge-weighted graph  $\mathcal{G} = (V(G) \cup X, E_1 \cup F_1 \cup F_2)$  from  $G_1$  as follows:

- $X = \{x_i, y_i, z_i \mid 1 \leq i \leq k\}$ .
- $F_1 = \{\{x_i, v\}, \{y_i, v\} \mid 1 \leq i \leq k, v \in V(C_i)\}$  and  $F_2 = \{\{x_i, z_i\}, \{y_i, z_i\} \mid 1 \leq i \leq k\}$ .
- The weight of each edge in  $E_1 \cup F_1$  is 0 while the weight of each edge in  $F_2$  is 1.
- For each  $v \in V(G)$ ,  $f(v) = g(v) = 2$ .
- For each  $i \in \{1, \dots, k\}$ ,  $f(x_i) = f(y_i) = f(z_i) = 0$ ,  $g(x_i) = g(y_i) = |V(C_i)|$ , and  $g(z_i) = 1$ .

For each weighted path-cycle cover  $M$  of  $G_1$ , we can obtain an  $[f, g]$ -factor  $N$  of  $\mathcal{G}$  from  $M$  as follows.

1. Initially,  $N = M$ .
2. For each  $i \in \{1, \dots, k\}$  and for each  $v \in V(C_i)$  with  $d_M(v) \leq 1$ , add edge  $\{v, x_i\}$  to  $N$  if  $d_M(v) = 1$ , and add edges  $\{v, x_i\}$  and  $\{v, y_i\}$  to  $N$  otherwise.
3. For each  $i \in \{1, \dots, k\}$  with  $d_N(x_i) < |V(C_i)|$  or  $d_N(y_i) < |V(C_i)|$ , add edge  $\{x_i, z_i\}$  to  $N$  if  $d_N(x_i) < |V(C_i)|$ , and add edge  $\{y_i, z_i\}$  to  $N$  otherwise.

Obviously, the weight of  $N$  is the same as that of  $M$ , i.e., equal to the number of odd cycles saturated by  $M$ . Thus, the maximum weight of an  $[f, g]$ -factor of  $\mathcal{G}$  is at least as large as the maximum weight of a path-cycle cover of  $G_1$ .

Conversely, for each maximum-weight  $[f, g]$ -factor  $N$  of  $\mathcal{G}$ , we can obtain a path-cycle cover  $M$  of  $G_1$  from  $N$  by letting  $E(M) = E(N) \cap E_1$ . We claim that the weight of  $M$  is the same as that of  $N$ . To see this claim, observe that for each  $i \in \{1, \dots, k\}$  with  $V(M) \cap V(C_i) \neq \emptyset$ ,

exactly one of edges  $\{x_i, z_i\}$  and  $\{y_i, z_i\}$  is contained in  $N$ . This observation holds because  $N$  is a maximum-weight  $[f, g]$ -factor of  $\mathcal{G}$ . By the claim, the maximum weight of a path-cycle cover of  $G_1$  is at least as large as the maximum weight of an  $[f, g]$ -factor of  $\mathcal{G}$ . So, by the discussion in the last paragraph, the maximum weight of a path-cycle cover of  $G_1$  is the same as the maximum weight of an  $[f, g]$ -factor of  $\mathcal{G}$ .  $\square$

Our algorithm then proceeds to perform the following four steps:

5. Compute a maximum-weight path-cycle cover  $M$  in  $G_1$ .
6. While there is an edge  $e \in M$  such that the weight of  $M - \{e\}$  is the same as that of  $M$ , delete  $e$  from  $M$ .
7. Construct a graph  $G_2 = (V(G), E(\mathcal{C}) \cup M)$ . (*Comment:* For each pair of connected components of  $\mathcal{C}$ , there is at most one edge between them in  $G_2$  because of Step 6.)
8. Construct a graph  $G_3$ , where the vertices of  $G_3$  one-to-one correspond to the connected components of  $\mathcal{C}$  and two vertices are adjacent in  $G_3$  if and only if the corresponding connected components of  $\mathcal{C}$  are adjacent in  $G_2$ .

**Fact 4.2** *Suppose that  $H$  is a connected component of  $G_3$ . Then, the following statements hold:*

1.  $H$  is a vertex, an edge, or a star.
2. If  $H$  is an edge, then at least one endpoint of  $H$  corresponds to an odd cycle of  $\mathcal{C}$ .
3. If  $H$  is a star, then every satellite of  $H$  corresponds to an odd cycle of  $\mathcal{C}$ .

An *isolated odd-cycle* of  $G_2$  is an odd cycle of  $G_2$  whose corresponding vertex in  $G_3$  is isolated in  $G_3$ . Similarly, a *leaf odd-cycle* of  $G_2$  is an odd cycle of  $G_2$  whose corresponding vertex in  $G_3$  is of degree 1 in  $G_3$ . Moreover, a *branching odd-cycle* of  $G_2$  is an odd cycle of  $G_2$  whose corresponding vertex in  $G_3$  is of degree 2 or more in  $G_3$ .

The next lemma is essentially the same as Lemma 2.1 in [9]. We include its proof here for self-containedness.

**Lemma 4.3** *Let  $I$  be the set of isolated odd-cycles in  $G_2$ . Then,  $|E(\mathcal{B})| \leq |E(\mathcal{C})| - |I|$ .*

PROOF. Let  $C_1, \dots, C_h$  be the odd cycles of  $\mathcal{C}$  such that for each  $i \in \{1, \dots, h\}$ ,  $\mathcal{B}$  contains no edge  $\{u, v\}$  with  $|\{u, v\} \cap V(C_i)| = 1$ . Let  $U_1 = \bigcup_{i=1}^h V(C_i)$  and  $U_2 = V(G) - U_1$ . For convenience, let  $C_0 = G[U_2]$ . Note that for each  $e \in E(\mathcal{B})$ , one of the graphs  $C_0, C_1, \dots, C_h$  contains both endpoints of  $e$ . So,  $\mathcal{B}$  can be partitioned into  $h + 1$  disjoint subgraphs  $\mathcal{B}_0, \dots, \mathcal{B}_h$  such that  $\mathcal{B}_i$  is a path-cycle cover of  $G[V(C_i)]$  for every  $i \in \{0, \dots, h\}$ . Since  $\mathcal{C}[U_2]$  must be a maximum-sized path-cycle cover of  $C_0$ ,  $|E(\mathcal{C}[U_2])| \geq |E(\mathcal{B}_0)|$ . The crucial point is that

for every  $i \in \{1, \dots, h\}$ ,  $|E(\mathcal{B}_i)| \leq |V(C_i)| - 1 = |E(C_i)| - 1$  because  $|V(C_i)|$  is odd. Thus,  $|E(\mathcal{C})| = |E(\mathcal{C}[U_2])| + \sum_{i=1}^h |E(C_i)| \geq |E(\mathcal{B}_0)| + \sum_{i=1}^h (|E(\mathcal{B}_i)| + 1) = |E(\mathcal{B})| + h$ .

Note that  $(V(G), E(G_1) \cap E(\mathcal{B}))$  is a path-cycle cover in  $G_1$  of weight  $k - h$ , where  $k$  is the number of odd cycles in  $\mathcal{C}$ . So,  $k - h \leq k - |I|$  because  $M$  is a maximum-weight path-cycle cover in  $G_1$  of weight  $k - |I|$ . So, by the last inequality in the last paragraph,  $|E(\mathcal{B})| \leq |E(\mathcal{C})| - h \leq |E(\mathcal{C})| - |I|$ .

□




Figure 1: An example of  $G_2$ , where the hollow vertices are free, the bold edges belong to  $\mathcal{C}$ , the left connected component is a bicycle, and the middle connected component is a tricycle.

Some definitions are in order (see Figure 1 for an example). A *bicycle* of  $G_2$  is a connected component of  $G_2$  that consists of two odd cycles and an edge between them. Note that a connected component of  $G_3$  is an edge if it corresponds to a bicycle in  $G_2$ . A *tricycle* of  $G_2$  is a connected component  $T$  of  $G_2$  that consists of one branching odd-cycle  $C_1$ , two leaf odd-cycles  $C_2$  and  $C_3$ , and two edges  $\{u_1, u_2\}$  and  $\{u_1, u_3\}$  such that  $u_1 \in V(C_1)$ ,  $u_2 \in V(C_2)$ , and  $u_3 \in V(C_3)$ . For convenience, we call  $C_1$  the *front cycle* of tricycle  $K$ , call  $C_2$  and  $C_3$  the *back cycles* of tricycle  $K$ , and call  $u_1$  the *front joint* of tricycle  $K$ .

A *cherry* of  $G_2$  is a subgraph  $Q$  of  $G_2$  that consists of two leaf odd-cycles  $C_1$  and  $C_2$  of  $\mathcal{C}$ , a vertex  $u \in V(G) - (V(C_1) \cup V(C_2))$ , and two edges  $\{u, v_1\}$  and  $\{u, v_2\}$  such that  $v_1 \in V(C_1)$  and  $v_2 \in V(C_2)$ . For convenience, we call edges  $\{u, v_1\}$  and  $\{u, v_2\}$  the *twigs* of cherry  $Q$ . By the construction of  $G_2$ , each pair of cherries are vertex-disjoint. Note that each odd cycle in a cherry of  $G_2$  is a satellite of a star in  $G_3$ . We classify the cherries of  $G_2$  into two types as follows. A cherry  $Q$  of  $G_2$  is of *type-1* if  $Q$  is a subgraph of a tricycle of  $G_2$ . Note that the two odd cycles in a type-1 cherry of  $G_2$  are the back cycles of a tricycle of  $G_2$ . A cherry of  $G_2$  is of *type-2* if it is not of type-1. Further note that there is no edge  $\{u, v\}$  in  $G$  such that  $u$  appears on an isolated odd-cycle of  $G_2$  and  $v$  appears on an odd cycle in a cherry of  $G_2$ .

A *lollipop* of  $G_2$  is a subgraph  $L$  of  $G_2$  that consists of a leaf odd-cycle  $C$  of  $G_2$ , a vertex  $u \notin V(C)$ , and an edge  $\{u, v\}$  with  $v \in V(C)$ . For convenience, we call edge  $\{u, v\}$  the *stick* of lollipop  $L$  and call vertex  $u$  the *end vertex* of lollipop  $L$ . A lollipop of  $G_2$  is *special* if it is neither a subgraph of a cherry of  $G_2$  nor a subgraph of a bicycle of  $G_2$ . A vertex  $u$  of  $G_2$  is *free* if no lollipop of  $G_2$  has  $u$  as its end vertex. Because of Step 3, each vertex of degree at most 2 in  $G_2$  is free.

We next define two types of operations that will be performed on  $G_2$ . An operation on  $G_2$  is *robust* if it removes no edge of  $\mathcal{C}$ , creates no new odd cycle, and creates no new isolated odd-cycle of  $G_2$ .






Figure 2: Three example cases of a type-1 operation, where the dotted lines are edges in  $E(G) - E(G_2)$ .

**Type 1:** Suppose that  $C$  is an odd cycle of a cherry  $Q$  of  $G_2$  and  $u$  is a free vertex of  $G_2$  with  $u \notin V(C)$  such that

- some vertex  $v$  of  $C$  is adjacent to  $u$  in  $G$  and
- if  $Q$  is a type-1 cherry of  $G_2$ , then  $u$  is not an endpoint of a twig of  $Q$ .

Then, a *type-1 operation* on  $G_2$  using cherry  $Q$  and edge  $\{u, v\}$  modifies  $G_2$  by performing the following steps (see Figure 2 for example cases):

- (1) If  $u$  appears on a leaf odd cycle  $C'$  of  $G_2$  such that  $C'$  is not part of a bicycle of  $G_2$  and  $Q$  is not a type-1 cherry of  $G_2$  with  $u \in V(Q)$ , then delete the stick of the lollipop containing  $C'$  from  $G_2$ .
- (2) Delete the twig of  $Q$  incident to a vertex of  $C$  from  $G_2$ .
- (3) Add edge  $\{u, v\}$  to  $G_2$ .

(*Comment:* A type-1 operation on  $G_2$  is robust and destroys at least one cherry of  $G_2$  without creating a new cherry in  $G_2$ .)

**Type 2:** Suppose that  $Q$  is a type-2 cherry of  $G_2$ ,  $B$  is a bicycle of  $G_2$ , and  $\{u, v\}$  is an edge in  $E(G_1) - E(G_2)$  such that  $u$  appears on an odd cycle  $C$  of  $Q$  and  $v$  appears on an odd cycle of  $B$ . Then, a *type-2 operation* on  $G_2$  using cherry  $Q$ , bicycle  $B$ , and edge  $\{u, v\}$  modifies  $G_2$  by deleting the twig of  $Q$  incident to a vertex of  $C$  and adding edge  $\{u, v\}$ .

(*Comment:* A type-2 operation on  $G_2$  is robust. Moreover, when no type-1 operation on  $G_2$  is possible, a type-2 operation on  $G_2$  destroys a type-2 cherry of  $G_2$  and creates a new type-1 cherry in  $G_2$ .)

Now, Step 9 of our algorithm is as follows.

9. While a type-1 or type-2 operation on  $G_2$  is possible, perform the following step:
- (a) If a type-1 operation on  $G_2$  is possible, perform a type-1 operation on  $G_2$ ; otherwise, perform a type-2 operation on  $G_2$ .

**Fact 4.4** *After Step 9, the following statements hold:*

1. *There is no edge  $\{u, v\}$  in  $E(G)$  such that  $u$  appears on an odd cycle in a type-2 cherry of  $G_2$  and  $v$  appears on another odd cycle in a type-2 cherry of  $G_2$ .*
2. *If  $\{u, v\}$  is an edge of  $G_1$  such that  $u$  appears on an odd cycle of a type-2 cherry of  $G_2$  and no type-2 cherry of  $G_2$  contains  $v$ , then  $v$  is the end vertex of a special lollipop or the front joint of a tricycle of  $G_2$ .*

Hereafter,  $G_2$  always refers to the graph obtained after the completion of Step 9.

Now, the final three steps of our algorithm are as follows:

10. Let  $U$  be the set of vertices that appear in type-2 cherries of  $G_2$ .
11. If  $U = \emptyset$ , then perform the following steps:
  - (a) For each connected component  $K$  of  $G_2$ , compute a maximum-sized edge-2-colorable subgraph of  $K$ . (*Comment:* Because of the simple structure of  $K$ , this step can be done in linear time by a standard dynamic programming.)
  - (b) Output the union of the edge-2-colorable subgraphs computed in Step 11a, and halt.
12. If  $U \neq \emptyset$ , then perform the following steps:
  - (a) Obtain an edge-2-colorable subgraph  $R$  of  $G - U$  by recursively calling the algorithm on  $G - U$ .
  - (b) For each type-2 cherry  $Q$  of  $G_2$ , obtain an edge-2-colorable subgraph of  $Q$  by removing one edge from each odd cycle  $C$  of  $Q$  that shares an endpoint with a twig of  $Q$ .
  - (c) Let  $\mathcal{A}_1$  be the union of  $R$  and the edge-2-colorable subgraphs computed in Step 12b.
  - (d) For each connected component  $K$  of  $G_2$ , compute a maximum-sized edge-2-colorable subgraph of  $K$ . (*Comment:* Because of the simple structure of  $K$ , this step can be done in linear time by a standard dynamic programming.)
  - (e) Let  $\mathcal{A}_2$  be the union of the edge-2-colorable subgraphs computed in Step 12d.
  - (f) If  $|E(\mathcal{A}_1)| \geq |E(\mathcal{A}_2)|$ , output  $\mathcal{A}_1$  and halt; otherwise, output  $\mathcal{A}_2$  and halt.

**Lemma 4.5** *Assume that  $G_2$  has no type-2 cherry. Then, the edge-2-colorable subgraph of  $G$  output in Step 11b contains at least  $r|E(\mathcal{B})|$  edges.*

PROOF. Let  $\mathcal{C}_2$  be the graph obtained from  $G_2$  by removing one edge from each isolated odd-cycle of  $G_2$ . By Lemma 4.3,  $|E(\mathcal{C}_2) \cap E(\mathcal{C})| \geq |E(\mathcal{B})|$ . Consider an arbitrary connected component  $K$  of  $\mathcal{C}_2$ . To prove the lemma, it suffices to prove that  $K$  has an edge-2-colorable subgraph  $K'$  with  $|E(K')| \geq r|E(K) \cap E(\mathcal{C})|$ . We distinguish several cases as follows:

*Case 1:*  $K$  is a bicycle of  $\mathcal{C}_2$ . To obtain an edge-2-colorable subgraph  $K'$  of  $K$ , we remove one edge  $e$  from each odd cycle of  $K$  such that one endpoint of  $e$  is of degree 3 in  $K$ . Note that  $|E(K')| = |E(K)| - 2 = |E(K) \cap E(\mathcal{C})| - 1$ . Since  $|E(K) \cap E(\mathcal{C})| \geq 10$ ,  $|E(K')| \geq \frac{9}{10}|E(K) \cap E(\mathcal{C})| > r|E(K) \cap E(\mathcal{C})|$ .

*Case 2:*  $K$  is a tricycle of  $\mathcal{C}_2$ . To obtain an edge-2-colorable subgraph  $K'$  of  $K$ , we first remove one edge  $e$  from each back odd-cycle of  $K$  such that one endpoint of  $e$  is of degree 3 in  $K$ , and then remove the two edges of the front odd-cycle incident to the vertex of degree 4 in  $K$ . Note that  $|E(K')| = |E(K)| - 4 = |E(K) \cap E(\mathcal{C})| - 2$ . Since  $|E(K) \cap E(\mathcal{C})| \geq 15$ ,  $|E(K')| \geq \frac{13}{15}|E(K) \cap E(\mathcal{C})| > r|E(K) \cap E(\mathcal{C})|$ .

*Case 3:*  $K$  is neither a bicycle nor a tricycle of  $\mathcal{C}_2$ . If  $K$  contains no odd cycle of  $\mathcal{C}$ , then  $K$  itself is edge-2-colorable and hence we are done. So, assume that  $K$  contains at least one odd cycle of  $\mathcal{C}$ . Then,  $K$  is also a connected component of  $G_2$ . Moreover, the connected component  $K''$  of  $G_3$  corresponding to  $K$  is either an edge or a star.

*Case 3.1:*  $K''$  is an edge. To obtain an edge-2-colorable subgraph  $K'$  of  $K$ , we start with  $K$ , delete the edge in  $E(K) - E(\mathcal{C})$ , and delete one edge from the unique odd cycle of  $K$ . Note that  $|E(K')| = |E(K)| - 2 = |E(K) \cap E(\mathcal{C})| - 1$ . Moreover,  $|E(K) \cap E(\mathcal{C})| \geq 7$  because of Step 3 and the robustness of Type-1 or Type-2 operations. Hence,  $|E(K')| \geq \frac{6}{7}|E(K) \cap E(\mathcal{C})| > r|E(K) \cap E(\mathcal{C})|$ .

*Case 3.2:*  $K''$  is a star. Let  $C_0$  be the connected component of  $\mathcal{C}$  corresponding to the center of  $K''$ . Let  $C_1, \dots, C_h$  be the odd cycles of  $\mathcal{C}$  corresponding to the satellites of  $K''$ . If  $C_0$  is a path, then  $K$  is a train graph and we are done by Lemma 3.1; otherwise,  $K$  is a starlike graph and we are done by Lemma 3.2.  $\square$

**Corollary 4.6** *If the maximum degree  $\Delta$  of a vertex in  $G$  is at most 3, then the ratio achieved by the algorithm is at least  $\frac{6}{7}$ .*

PROOF. When  $\Delta \leq 3$ ,  $G_2$  has no cherry because of Step 3. Moreover, Lemmas 3.1, 3.2, and 4.5 still hold even when we replace the ratio  $r$  by  $\frac{6}{7}$ .  $\square$

In order to analyze the approximation ratio achieved by our algorithm when  $G_2$  has at least one type-2 cherry after Step 9, we need to define several notations as follows:

- Let  $s$  be the number of special lollipops in  $G_2$ .
- Let  $t$  be the number of tricycles in  $G_2$ .

- Let  $c$  be the number of type-2 cherries in  $G_2$ .
- Let  $\ell$  be the total number of vertices that appear on odd cycles in the type-2 cherries in  $G_2$ .

**Lemma 4.7** *Let  $E(\mathcal{B}_2)$  be the set of all edges  $e \in E(\mathcal{B})$  such that at least one endpoint of  $e$  appears in a type-2 cherry of  $G_2$ . Then,  $|E(\mathcal{B}_2)| \leq \ell + 2s + 2t$ .*

PROOF.  $E(\mathcal{B}_2)$  can be partitioned into the following three subsets:

- $E(\mathcal{B}_{2,1})$  consists of those edges  $e \in E(\mathcal{B})$  such that at least one endpoint of  $e$  is the vertex of a type-2 cherry of  $G_2$  that is a common endpoint of the two twigs of the cherry.
- $E(\mathcal{B}_{2,2})$  consists of those edges  $e \in E(\mathcal{B})$  such that each endpoint of  $e$  appears on an odd cycle of a type-2 cherry of  $G_2$ .
- $E(\mathcal{B}_{2,3})$  consists of those edges  $\{u, v\} \in E(\mathcal{B})$  such that  $u$  appears on an odd cycle of a type-2 cherry of  $G_2$  and no type-2 cherry of  $G_2$  contains  $v$ .

Obviously,  $|E(\mathcal{B}_{2,1})| \leq 2c$ . By Statement 1 in Fact 4.4,  $|E(\mathcal{B}_{2,2})| \leq \ell - 2c$  because for each odd cycle  $C$ ,  $\mathcal{B}_{2,2}$  can contain at most  $|V(C)| - 1$  edges  $\{u, v\}$  with  $\{u, v\} \subseteq V(C)$ . By Statement 2 in Fact 4.4,  $|E(\mathcal{B}_{2,3})| \leq 2s + 2t$ . So,  $|E(\mathcal{B}_2)| \leq \ell + 2s + 2t$ .  $\square$

**Lemma 4.8** *The ratio achieved by the algorithm is at least  $r$ .*

PROOF. The proof is by induction on  $|V(G)|$ , the number of vertices in the input graph  $G$ . If  $|V(G)| \leq 2$ , then our algorithm outputs a maximum-sized edge-2-colorable subgraph of  $G$ . So, assume that  $|V(G)| \geq 3$ . Then, after our algorithm finishes executing Step 10, the set  $U$  may be empty or not. If  $U = \emptyset$ , then by Lemma 4.5, the edge-2-colorable subgraph output by our algorithm has at least  $r|E(\mathcal{B})|$  edges and we are done. So, suppose that  $U \neq \emptyset$ .

First consider the case where  $s + t \leq \frac{1-r}{2r}\ell$ . In this case,  $\frac{\ell+r|E(\mathcal{B}_1)|}{\ell+2s+2t+|E(\mathcal{B}_1)|} \geq r$ , where  $\mathcal{B}_1$  is a maximum-sized edge-2-colorable subgraph of  $G - U$ . Moreover, by the inductive hypothesis,  $|E(\mathcal{A}_1)| \geq \ell + r|E(\mathcal{B}_1)|$ . Furthermore, by Lemma 4.7,  $|E(\mathcal{B})| \leq \ell + 2s + 2t + |E(\mathcal{B}_1)|$ . So, the lemma holds in this case.

Next consider the case where  $s + t > \frac{1-r}{2r}\ell$ . Let  $\mathcal{C}_2$  be the graph obtained from  $G_2$  by removing one edge from each isolated odd-cycle of  $G_2$ . By Lemma 4.3,  $|E(\mathcal{C}_2) \cap E(\mathcal{C})| \geq |E(\mathcal{B})|$ . Let  $\mathcal{C}_3$  be the graph obtained from  $\mathcal{C}_2$  by removing one twig from each type-2 cherry. Note that there are exactly  $c$  isolated odd-cycles in  $\mathcal{C}_3$ . Moreover, since the removed twig does not belong to  $E(\mathcal{C})$ , we have  $|E(\mathcal{C}_3) \cap E(\mathcal{C})| \geq |E(\mathcal{B})|$ . Consider an arbitrary connected component  $K$  of  $\mathcal{C}_3$ . To prove the lemma, we want to prove that  $K$  has an edge-2-colorable subgraph  $K'$  with  $|E(K')| \geq r|E(K) \cap E(\mathcal{C})|$ . This goal can be achieved because of Lemma 4.5, when  $K$  is not an isolated odd-cycle. On the other hand, this goal can not be achieved when  $K$  is an isolated odd-cycle (of length at least 5). Our idea behind the proof is to charge the deficit in the edge numbers of isolated odd-cycles of  $\mathcal{C}_3$  to the other connected components of  $K$  because they have surplus in their edge numbers.

The deficit in the edge number of each isolated odd-cycle of  $\mathcal{C}_3$  is at most  $5r - 4$ . So, the total deficit in the edge numbers of the isolated odd-cycles of  $\mathcal{C}_3$  is at most  $(5r - 4)c$ . We charge a penalty of  $6 - 7r$  to each non-isolated odd-cycle of  $\mathcal{C}_3$  that is also an odd cycle in a type-2 cherry of  $G_2$  or is also the odd cycle in a special lollipop of  $G_2$ . We also charge a penalty of  $\frac{6-7r}{3}$  to each odd cycle of  $\mathcal{C}_3$  that is part of a tricycle of  $G_2$ . Clearly, the total penalties are  $(6 - 7r)c + (6 - 7r)(s + t) > (6 - 7r)c + \frac{(6-7r)(1-r)}{2r}\ell$ . Note that  $\ell \geq 10c$ . The total penalties are thus at least  $(6 - 7r)c + \frac{5(6-7r)(1-r)}{r}c = \frac{30-59r+28r^2}{r}c \geq (5r - 4)c$ , where the last inequality follows from the equation  $23r^2 - 55r + 30 = 0$ . So, the total penalties are at least as large as the total deficit in the edge numbers of the isolated odd-cycles of  $\mathcal{C}_3$ . Therefore, to prove the lemma, it suffices to prove that for every connected component  $K$  of  $\mathcal{C}_3$ , we can compute an edge-2-colorable subgraph  $K'$  of  $K$  such that  $|E(K')| - p(K) \geq r|E(K) \cap E(\mathcal{C})|$ , where  $p(K)$  is the total penalties of the odd cycles in  $K$ . As in the proof of Lemma 4.5, we distinguish several cases as follows:

*Case 1:*  $K$  is a bicycle of  $\mathcal{C}_2$ . In this case,  $p(K) = 0$ . Moreover, we can compute an edge-2-colorable subgraph  $K'$  of  $K$  such that  $|E(K')| \geq \frac{9}{10}|E(K) \cap E(\mathcal{C})|$  (cf. Case 1 in the proof of Lemma 4.5). So,  $|E(K')| - p(K) \geq r|E(K) \cap E(\mathcal{C})|$  because  $r \leq \frac{9}{10}$ .

*Case 2:*  $K$  is a tricycle of  $\mathcal{C}_2$ . In this case,  $p(K) = 6 - 7r$ . Moreover, we can compute an edge-2-colorable subgraph  $K'$  of  $K$  such that  $|E(K')| = |E(K) \cap E(\mathcal{C})| - 2$  (cf. Case 2 in the proof of Lemma 4.5). So,  $|E(K')| - p(K) \geq r|E(K) \cap E(\mathcal{C})|$  because  $|E(K) \cap E(\mathcal{C})| \geq 15$  and  $r \leq \frac{7}{8}$ .

*Case 3:*  $K$  is neither a bicycle nor a tricycle of  $\mathcal{C}_2$ . We may assume that  $K$  contains at least one odd cycle of  $\mathcal{C}$ . Then,  $K$  is also a connected component of  $G_2$ . Moreover, the connected component  $K''$  of  $G_3$  corresponding to  $K$  is either an edge or a star.

*Case 3.1:*  $K''$  is an edge. In this case,  $p(K) \leq 6 - 7r$ . Moreover, we can compute an edge-2-colorable subgraph  $K'$  of  $K$  such that  $|E(K')| = |E(K) \cap E(\mathcal{C})| - 1$  (cf. Case 3.1 in the proof of Lemma 4.5). So,  $|E(K')| - p(K) \geq r|E(K) \cap E(\mathcal{C})|$  because  $|E(K) \cap E(\mathcal{C})| \geq 7$ .

*Case 3.2:*  $K''$  is a star. Let  $C_0$  be the connected component of  $\mathcal{C}$  corresponding to the center of  $K''$ . Let  $C_1, \dots, C_h$  be the odd cycles of  $\mathcal{C}$  corresponding to the satellites of  $K''$ . If  $C_0$  is a path, then  $K$  is a train graph and we are done by Lemma 3.1; otherwise,  $K$  is a starlike graph and we are done by Lemma 3.2.  $\square$

Clearly, each step of our algorithm except Step 12a can be implemented in  $O(n^2m)$  time. Since the recursion depth of the algorithm is  $O(n)$ , it runs in  $O(n^3m)$  total time. In summary, we have shown the following theorem:

**Theorem 4.9** *There is an  $O(n^3m)$ -time approximation algorithm for MAX SIMPLE EDGE 2-COLORING that achieves a ratio of roughly 0.842.*

## 5 An Application

Let  $G$  be a graph. An *edge cover* of  $G$  is a set  $F$  of edges of  $G$  such that each vertex of  $G$  is incident to at least one edge of  $F$ . For a natural number  $k$ , a  $[1, \Delta]$ -factor  $k$ -packing of  $G$  is a collection of  $k$  disjoint edge covers of  $G$ . The *size* of a  $[1, \Delta]$ -factor  $k$ -packing  $\{F_1, \dots, F_k\}$  of  $G$  is

$|F_1| + \dots + |F_k|$ . The problem of deciding whether a given graph has a  $[1, \Delta]$ -factor  $k$ -packing was considered in [7, 8]. In [10], Kosowski *et al.* defined the *minimum  $[1, \Delta]$ -factor  $k$ -packing problem* (MIN- $k$ -FP) as follows: Given a graph  $G$ , find a  $[1, \Delta]$ -factor  $k$ -packing of  $G$  of minimum size or decide that  $G$  has no  $[1, \Delta]$ -factor  $k$ -packing at all.

According to [10], MIN-2-FP is of special interest because it can be used to solve a fault tolerant variant of the guards problem in grids (which is one of the art gallery problems [11, 12]). Indeed, they proved the NP-hardness of MIN-2-FP and the following lemma:

**Lemma 5.1** *If MAX SIMPLE EDGE 2-COLORING admits an approximation algorithm  $A$  achieving a ratio of  $\alpha$ , then MIN-2-FP admits an approximation algorithm  $B$  achieving a ratio of  $2 - \alpha$ . Moreover, if the time complexity of  $A$  is  $T(n)$ , then the time complexity of  $B$  is  $O(T(n))$ .*

So, by Theorem 4.9, we have the following immediately:

**Theorem 5.2** *There is an  $O(n^3m)$ -time approximation algorithm for MIN-2-FP achieving a ratio of roughly 1.158.*

## 6 Open Problems

One obvious open question is to ask whether one can design a polynomial-time approximation algorithm for MAX SIMPLE EDGE 2-COLORING that achieves a ratio significantly better than 0.842. Assuming  $P \neq NP$ , the APX-hardness proof of the problem given in [3] implies a lower bound of roughly 0.999937 on the ratio achievable by a polynomial-time approximation algorithm. It seems interesting to prove a significantly better lower bound.

## Acknowledgments

We thank the referees for helpful comments.

## References

- [1] Z.-Z. Chen and R. Tanahashi. Approximating Maximum Edge 2-Coloring in Simple Graphs via Local Improvement. *Theoretical Computer Science* (special issue on AAIM 2008), **410** (2009) 4543-4553. A preliminary version appeared in *Proceedings of 4th International Conference on Algorithmic Aspects in Information and Management* (AAIM 2008), Lecture Notes in Computer Science, **5034** (2008) 84-96.
- [2] Z.-Z. Chen, R. Tanahashi, and L. Wang. An Improved Approximation Algorithm for Maximum Edge 2-Coloring in Simple Graphs. *Journal of Discrete Algorithms*, Vol. 6, No. 2, pp. 205-215, 2008. A preliminary version appeared in *Proceedings of 3rd International Conference on Algorithmic Aspects in Information and Management*, Lecture Notes in Computer Science, **4508** (2007) 27-36.

- [3] U. Feige, E. Ofek, and U. Wieder. Approximating Maximum Edge Coloring in Multigraphs. *Proceedings of the 10th International Conference on Integer Programming and Combinatorial Optimization (IPCO)*, Lecture Notes in Computer Science, **2462** (2002) 108-121.
- [4] H. Gabow. An Efficient Reduction Technique for Degree-Constrained Subgraph and Bidirected Network Flow Problems. *Proceedings of the 15th Annual ACM Symposium on Theory of Computing (STOC'83)*, pp. 448-456, ACM, 1983.
- [5] D. Hartvigsen. *Extensions of Matching Theory*. Ph.D. Thesis, Carnegie-Mellon University, 1984.
- [6] D. Hochbaum. *Approximation Algorithms for NP-Hard Problems*. PWS Publishing Company, Boston, 1997.
- [7] D.P. Jacobs and R.E. Jamison. Complexity of Recognizing Equal Unions in Families of Sets. *Journal of Algorithms*, **37** (2000) 495-504.
- [8] K. Kawarabayashi, H. Matsuda, Y. Oda, and K. Ota. Path Factors in Cubic Graphs. *Journal of Graph Theory*, **39** (2002) 188-193.
- [9] A. Kosowski. Approximating the Maximum 2- and 3-Edge-Colorable Subgraph Problems. *Discrete Applied Mathematics*, **157** (2009) 3593-3600.
- [10] A. Kosowski, M. Malafiejski, and P. Zylinski. Packing  $[1, \Delta]$ -Factors in Graphs of Small Degree. *Journal of Combinatorial Optimization*, **14** (2007) 63-86.
- [11] J. O'Rourke. *Art Gallery Theorems and Algorithms*. Oxford University Press, 1987.
- [12] J. Urrutia. *Art Gallery and Illumination Problems*. Handbook on Computational Geometry, Elsevier Science, Amsterdam, 2000.